



लक्ष्मी सीड्स

Suggestions For Higher Production





लक्ष्मी सीड्स

Research Guar Varieties

Every seed genuine, brimming with prosperity.

Aakash Hara Sona Mukhiya



Field Preparation and Soil Type

Sowing can be done in all types of soil; it requires one or two deep ploughings and leveling.



Time of sowing

For good production, the sowing of cluster bean (guar) is suitable from June 15 to July 30.



Seed quantity

2 kg per acre for rain-fed and monsoon-sown areas
seed rate of 3 kg per acre.



Seed Treatment and Sowing

This seed is treated; you can treat it with
“*Trichoderma viride*”.



Quantity of manure and fertilizers

Mix 25–30 kg of urea and 120–130 kg of single superphosphate per acre into the field before sowing.



Irrigation

1–2 irrigations (in irrigated areas); dependence solely on rainfall in unirrigated areas.



Weed Control

Dissolve 750 ml of Stomp 30% EC (Pendimethalin) in 200 liters of water and spray it per acre using a flood jet nozzle within 24 hours of sowing.



Harvesting

The pods of this variety do not shatter or drop, allowing for simultaneous harvesting.



Comment

Note: The advice provided above is based on our experience, field trials, and experiments conducted at research centers. However, variations may occur due to natural factors such as climate, soil type, and farming practices. We cannot provide guarantees beyond seed quality, as other conditions are beyond our control. The information provided here is intended for the prosperity and success of our fellow farmers.

